

Early Diagnosis of COPD does help!

COPD (Chronic Obstructive Pulmonary Disease) is an increasingly common condition resulting in considerable morbidity and mortality. This opinion sheet will review the evidence for the benefits of early diagnosis and give you some tools that you can use to make changes in your practice to improve health outcomes for your patients.

Why does early diagnosis matter?

COPD is commonly under-diagnosed worldwide. Only a quarter of the people shown to have COPD in a population survey in Spain were previously aware of the diagnosis.¹

Diagnosis of COPD usually does not occur until significant lung function has already been lost. By the time patients recognise that they have symptoms, their FEV₁ has usually fallen to about 50% predicted - a level where health status is already reduced and there is a significant amount of systemic inflammation leading to co-morbidities.²

Financial costs of COPD are high. These include the direct costs of hospitalisation and other healthcare interventions, as well as the indirect costs of disability, lost productivity, caregiver support and family costs. Many of these costs could be reduced by earlier diagnosis and intervention.³

Can early intervention help?

At all stages in the management of COPD, interventions including smoking cessation, exercise and rehabilitation, lifestyle changes, influenza and pneumococcal vaccination and the reduction of exacerbations result in better quality of life for the patient.²

A growing body of evidence suggests that early detection of airflow limitation and early intervention can delay lung function decline, reduce the burden of COPD symptoms, and improve patients' quality of life.⁴

Smoking cessation has been shown to be the

most significant intervention to slow the rate of decline of lung function⁵ and the earlier a smoker stops smoking, the better lung function is preserved.⁶ Smoking cessation intervention may well be more successful in those who are actually given a firm diagnosis.⁷ Reviewing the illness and instructing patients on their lung age has also been shown to improve smoking cessation rates.⁸ For tools on smoking cessation please see the IPCRG smoking cessation fact sheet⁹ or the web-based practical guidance.¹⁰

Early pharmacological intervention can improve the health status and exercise capacity of COPD patients, and reduce exacerbations, even in patients with mild to moderate COPD.¹¹ Airflow limitation during exercise is associated with extensive small airways dysfunction even in patients whose lung function at rest may appear to be relatively preserved. These patients are likely to benefit from bronchodilation irrespective of the improvements observed with spirometry.¹² Early identification also allows earlier lifestyle change such as exercise and pulmonary rehabilitation.¹³

Early and aggressive management of exacerbations protects the patient from COPD progression. Every purulent COPD exacerbation decreases quality of life and longevity, and reduces lung function.¹⁴

Vaccination, adequate nutrition, and appropriate pharmacological intervention have been shown to decrease exacerbations.^{15,16}

Self-management education is a crucial component of care and all people with COPD should be offered the opportunity to discuss the lifestyle changes that can improve prognosis and develop plans for early intervention of exacerbations.¹⁷

What are the barriers to making the diagnosis earlier?

There are many barriers to an early diagnosis - see Table 1.

How do we promote earlier diagnosis?

There are a number of strategies that can be used to encourage earlier diagnosis. Promoting better understanding and awareness among politicians, health professionals and people in

Table 1. Barriers when promoting COPD earlier diagnosis

- COPD progresses relentlessly, but slowly, and as such, many patients do not realise that they have a problem
- COPD patients blame their breathlessness on aging, being less active and becoming older. They assume their cough is a normal phenomenon; the 'smoker's cough'
- COPD patients tend to be uncomplaining about their condition – described as the 'silence of people with COPD'¹⁸
- Because patients under-emphasise their symptoms, the physician may be less aggressive about treating them and does not consider the disease at an early stage
- Physicians might not consider repeated bronchial infections as an early sign of COPD development
- These patients often have multiple co-morbidities, and these conditions may well be more pressing and clearer to diagnose
- Physicians have a gender bias, assuming that females would have asthma, and thus missing the proper diagnosis¹⁹
- Controversy about the use of spirometry in primary care for early detection may discourage some clinicians²⁰
- There is a lack of consistently performed spirometry and spirometry training in primary care²¹
- There may be a delay in receiving spirometry reports when done outside of the office²²
- There are time pressures on General Practitioners that impact adversely on their capacity to manage patients proactively^{23,24}

the community about an increasingly important disease is the cornerstone of a global change in attitude.


Offering spirometry to all smokers regardless of whether they are symptomatic for COPD has been advocated, with a detection rate for newly diagnosed COPD of up to 20%.²⁵ However, many authors recommend case finding by offering spirometry to symptomatic smokers after an initial approach using existing questionnaires to detect COPD-related initial symptoms.²⁶

In a population setting, questionnaires could be distributed through the media to encourage smokers at risk to check whether they have symptoms and visit their general practitioners. The Canada Lung Health Test²⁷ is one good example (Table 2). A negative screen makes a diagnosis of COPD less likely.²⁸

The IPCRG currently recommends that all patients over 35 years old should be evaluated for their risk of developing COPD²⁹ - see Figure 1.

1. Take a history using validated screening questionnaires based on a combination of risk factors and symptoms.
2. Perform 'case-identification' spirometry using a variety of small "mini-spirometers" that can exclude those with normal FEV₁ and identify those who require more complete investigation for COPD.
3. Offer diagnostic spirometry to patients who have either symptoms and risk factors, or a positive screening questionnaire, or whose screening FEV₁ is not within normal limits.

Figure 1.²⁹ An approach to COPD case-identification in primary care
Adapted with permission from *Primary Care Respiratory Journal*
Full text of this article is available at: <http://dx.doi.org/10.4104/pcrj.2009.00055>



Concern has been expressed about the accuracy of spirometry performed in primary care settings^{30,31} However, studies show that accurate spirometry can be performed in primary care offices, where the operators have appropriate training and interest.

Practices that have introduced spirometry into

routine care have made significant changes in COPD diagnosis and treatment.³²

Opportunistic spirometry to detect COPD has been shown to be cost effective.³³

Summary

Early diagnosis of patients with COPD is good for the patient and the community. The health care worker must suspect the possible diagnosis from symptoms and risk factors, consider screening with mini-spirometers, and offer proper spirometry to confirm the diagnosis. Making the diagnosis early will encourage smoking cessation and enable earlier interventions to help prevent exacerbations and hopefully preserve lung function, quality of life and decrease mortality.

Table 2. Canada Lung Health Test²⁷

1. Do you cough regularly?
2. Do you cough up phlegm regularly?
3. Do even simple chores make you short of breath?
4. Do you wheeze when you exert yourself, or at night?
5. Do you get frequent colds that persist longer than those of other people you know?

If the patient is a smoker or ex-smoker and over 40 years old and answers yes to any of the listed questions, referral should be made for further assessment, including spirometry.

References

1. Miravittles M, Soriano JB, Garcia-Rio F, et al. Prevalence of COPD in Spain: impact of undiagnosed COPD on quality of life and daily life activities *Thorax*. 2009;**64**(10):863-8. Epub 2009 Jun 23.
2. Global Strategy for the Diagnosis, Management and Prevention of COPD. Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2008. Available from: <http://www.goldcopd.org>.
3. Tzovaras NZ, Koulourmanis VN, Gourgoulis KI. The economic impact of late detection of COPD in general practice. *Chest* 2005;**127**(1):412.
4. Tinkelman DG, Price D, Nordyke RJ, Halbert RJ. COPD screening efforts in primary care: what is the yield? *Prim Care Resp J* 2007;**16**(1):41-8.
5. Fletcher C, Peto R. A prospective epidemiological study of the natural history of chronic airflow obstruction. *Br Med J*. 1977;**25**;1:1645-8.
6. Scanlon PD, Connett JE, Waller LA, et al. Smoking cessation and lung function in mild-to-moderate chronic obstructive pulmonary disease. The Lung Health Study. *Am J Respir Crit Care Med* 2000;**161**(2 Pt 1):381-90.
7. Bednarek M, Gorecka D, Wielgomas J, et al. Smokers with airway obstruction are more likely to quit smoking. *Thorax* 2006;**61**(10):869.
8. Parkes G, Greenhalgh T, Griffin M, et al. Effect on smoking quit rate of telling patients their lung age: the Step2quit randomised controlled trial. *BMJ* 2008;**336**:598-600.

9. van Schayck OCP, Pinnock H, Ostrem A, Litt J for the IPCRG. IPCRG Consensus statement: Tackling the smoking epidemic - practical guidance for primary care. *Prim Care Respir J* 2008;**17**(3): 185-193 DOI: <http://dx.doi.org/10.3132/pcrj.2008.00060>
10. IPCRG web-based practical guidance <http://www.theipcr.org/smoking/index.php>
11. Johansson G, Lindberg A, Romberg K, et al. Bronchodilator efficacy of tiotropium in patients with mild to moderate COPD. *Prim Care Resp J* 2008;**17**(3):169-175.
12. Ofir D, Laveneziana P, Webb KA, et al. Mechanisms of dyspnea during cycle exercise in symptomatic patients with GOLD stage I chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2008;**177**:622-9.
13. Lacasse Y, Goldstein R, Lasserson TJ, Martin S. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2006;(4):CD003793. DOI: <http://dx.doi.org/10.1002/14651858.CD003793.pub2>.
14. . JJ. Soler-Cataluna, M.A. Martinez-Garcia, P. Roman Sanchez, E. et al. Severe acute exacerbations and mortality in patients with chronic obstructive pulmonary disease: *Thorax* 2005;**60**:925-931.
15. Calverley PMA, Anderson JA, Celli B et al. for the TORCH investigators Salmeterol and Fluticasone Propionate and Survival in Chronic Obstructive Pulmonary Disease. *N Engl J Med* 2007;**356**:775-789.
16. Tashkin DP, Celli B, Senn S, et al. A 4-Year Trial of Tiotropium in Chronic Obstructive Pulmonary Disease. *N Engl J Med* 2008;**359**:1543-54.
17. Kaplan A. The COPD Action Plan. *Can Fam Physician*. 2009;**55**(1):58-9.
18. Habraken JM, Pols J, Bindels PJE, Willems DL. The silence of patients with end-stage COPD: a qualitative study. *Br J Gen Pract* 2008;**58**:844-849.
19. Chapman KR, Tashkin, DP, Pye D. Gender Bias in the Diagnosis of COPD. *Chest*. 2001;**119**:1691-1695.
20. Lin K, Watkins B, Johnson T, et al. Screening for Chronic Obstructive Pulmonary Disease Using Spirometry: Summary of the Evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2008;**148**:60520-213.

21. Lee TA, Bartle B, Weiss KB. Spirometry use in clinical practice following diagnosis of COPD. *Chest* 2006;**129**:1509-1515
22. Stanbrook MB, Kaplan A. The error of not measuring asthma. *CMAJ*. 2008;**179**:1099-102
23. Østbye T, Yarnall KSH, Krause KM, et al. Is There Time for Management of Patients With Chronic Diseases in Primary Care? *Ann Fam Med* 2005;**3**:209-214.
24. Yarnall KS, Østbye T, Krause KM, P et al. Family physicians as team leaders: "time" to share the care. *Prev Chronic Dis*. 2009;**6**(2):A59. Epub 2009 Mar 16.
26. Qaseem A, Snow V, Shekelle P, et al. Diagnosis and management of stable chronic obstructive pulmonary disease: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2007;**147**(9):633-8.
27. O'Donnell D, Hernandez P, Kaplan A, et al. Canadian Thoracic Society recommendations for management of chronic obstructive lung disease-2008 update- highlights for primary care. *Can Resp J* 2008;**15**(Suppl A): 1A-8A
28. Jing J-Y, Huang TC, Cui W, et al. Should FEV₁/FEV₆ replace FEV₁/FVC ratio to detect airway obstruction? A meta-analysis. *Chest* 2009;**135**(4):991-8.
29. Price D, Crockett A, Arne M, et al. Spirometry in primary care case-identification, diagnosis and management of COPD. *Prim Care Respir J* 2009; **18**(3):216-223. DOI: <http://dx.doi.org/10.4104/pcrj.2009.00055>
30. Enright P. Provide GPs with spirometry, not spirometers. *Thorax* 2008;**63**(5):387 and
31. Soriano JB, Zielinski J, Price D. Screening for and early detection of chronic obstructive pulmonary disease. *Lancet*. 2009;**374**:721-32.
32. Miravittles M, de la Roza C, Naberan K, et al. Use of spirometry and patterns of prescribing in COPD in primary care. *Respir Med* 2007;**101**(8):1753.
33. National Collaborating Centre for Chronic Conditions. Chronic obstructive pulmonary disease. Management of chronic obstructive pulmonary disease in adults in primary and secondary care. National Institute for Clinical Excellence 2004.