

## Breathlessness in adults: A practical guide for primary care clinicians

### References

- Poulos LM, Ampon RD, Currow DC, Marks GB, Toelle BG, Reddel HK. Prevalence and burden of breathlessness in Australian adults: The National Breathlessness Survey—a cross-sectional web-based population survey. *Respirology*. 2021.
- Ekström M, Sundh J, Andersson A, Angerås O, Blomberg A, Börjesson M, et al. Exertional breathlessness related to medical conditions in middle-aged people: the population-based SCAPIS study of more than 25,000 men and women. *Respiratory Research*. 2024;25(1):127.
- Sandberg J, Ekström M, Börjesson M, Bergström G, Rosengren A, Angerås O, Toren K. Underlying contributing conditions to breathlessness among middle-aged individuals in the general population: a cross-sectional study. *BMJ Open Respir Res*. 2020;7(1).
- Santos M, Kitzman DW, Matsushita K, Loehr L, Sueta CA, Shah AM. Prognostic Importance of Dyspnea for Cardiovascular Outcomes and Mortality in Persons without Prevalent Cardiopulmonary Disease: The Atherosclerosis Risk in Communities Study. *PLoS one*. 2016;11(10):e0165111.
- Nishimura K, Izumi T, Tsukino M, Oga T. Dyspnea is a better predictor of 5-year survival than airway obstruction in patients with COPD. *Chest*. 2002;121(5):1434-40.
- Sandberg J, Engström G, Ekström M. Breathlessness and incidence of COPD, cardiac events and all-cause mortality: A 44-year follow-up from middle age throughout life. *PLoS one*. 2019;14(3):e0214083.
- Sunjaya A, Martin A, Arnott C, Jenkins C. Management of chronic breathlessness in primary care: what do GPs, non-GP specialists, and allied health professionals think? *Aust J Prim Health*. 2023;29(4):375-84.
- Doe G, Clanchy J, Wathall S, Chantrell S, Edwards S, Baxter N, et al. Feasibility study of a multicentre cluster randomised control trial to investigate the clinical and cost-effectiveness of a structured diagnostic pathway in primary care for chronic breathlessness: protocol paper. *BMJ Open*. 2021;11(11):e057362.
- Doe G, Clanchy J, Wathall S, Barber S, Edwards SA, Evans H, Jackson D, Armstrong N, Steiner MC, Evans RA. Investigating a structured diagnostic approach for chronic breathlessness in primary care: a mixed-methods feasibility cluster Randomised Controlled Trial *medRxiv* 2024.02.21.24303140
- Bausewein C, Booth S, Gysels M, Higginson I. Non-pharmacological interventions for breathlessness in advanced stages of malignant and non-malignant diseases. *Cochrane Database Syst Rev*. 2008(2):Cd005623.
- Parshall MB, Schwartzstein RM, Adams L, Banzett RB, Manning HL, Bourbeau J, et al. An official American Thoracic Society statement: update on the mechanisms, assessment, and management of dyspnea. *American journal of respiratory and critical care medicine*. 2012;185(4):435-52.
- Johnson MJ, Yorke J, Hansen-Flaschen J, Lansing R, Ekstrom M, Similowski T, Currow DC. Towards an expert consensus to delineate a clinical syndrome of chronic breathlessness. *The European respiratory journal*. 2017;49(5).
- Currow DC, Dal Grande E, Ferreira D, Johnson MJ, McCaffrey N, Ekstrom M. Chronic breathlessness associated with poorer physical and mental health-related quality of life (SF-12) across all adult age groups. *Thorax*. 2017;72(12):1151-3.
- Ramon MA, Ter Riet G, Carsin AE, Gimeno-Santos E, Agustí A, Antó JM, et al. The dyspnoea-inactivity vicious circle in COPD: development and external validation of a conceptual model. *The European respiratory journal*. 2018;52(3).
- Murtagh GM. A critical look at ideas, concerns and expectations in clinical communication. *Med Educ*. 2023;57(4):331-6.
- Lunn S, Dharmagunawardena R, Lander M, Sweeney J. It's hard to talk about breathlessness: a unique insight from respiratory trainees. *Clin Med (Lond)*. 2019 Jul;19(4):344-347. doi: 10.7861/clinmedicine.19-4-344
- NHS England. Adult breathlessness pathway (pre-diagnosis): diagnostic pathway support tool; <https://www.england.nhs.uk/long-read/adult-breathlessness-pathway-pre-diagnosis-diagnostic-pathway-support-tool/> 2024 Accessed: 2024-04-02.
- Baxter N. Breathlessness in the primary care setting. *Current Opinion in Supportive & Palliative Care*. 2017;11(3):152-8.
- Mahler DA, Wells CK. Evaluation of clinical methods for rating dyspnea. *Chest*. 1988;93(3):580-6.
- van Dixhoorn J, Folgering H. The Nijmegen Questionnaire and dysfunctional breathing. *ERJ Open Res*. 2015;1(1).
- Spathis A, Booth S, Moffat C, Hurst R, Ryan R, Chin C, Burkin J. The Breathing, Thinking, Functioning clinical model: a proposal to facilitate evidence-based breathlessness management in chronic respiratory disease. *NPJ Prim Care Respir Med*. 2017;27(1):27.
- Singh B, Olds T, Curtis R, Dumuid D, Virgara R, Watson A, Szeto K, O'Connor E, Ferguson T, Eglitis E, Miatke A, Simpson CE, Maher C. Effectiveness of physical activity interventions for improving depression, anxiety and distress: an overview of systematic reviews. *Br J Sports Med*. 2023 Sep;57(18):1203-1209. doi: 10.1136/bjsports-2022-106195
- Semlitsch T, Stigler FL, Jeitler K, Horvath K, Siebenhofer A. Management of overweight and obesity in primary care—A systematic overview of international evidence-based guidelines. *Obes Rev*. 2019;20(9):1218-30.
- Ekström M, Ferreira D, Chang S, Louw S, Johnson MJ, Eckert DJ, et al. Effect of Regular, Low-Dose, Extended-release Morphine on Chronic Breathlessness in Chronic Obstructive Pulmonary Disease: The BEAMS Randomized Clinical Trial. *Jama*. 2022;328(20):2022-32.
- Currow DC, Ekström M, Louw S, Hill J, Fazekas B, Clark K, et al. Sertraline in symptomatic chronic breathlessness: a double blind, randomised trial. *The European respiratory journal*. 2019;53(1).
- Simon ST, Higginson IJ, Booth S, Harding R, Weingartner V, Bausewein C. Benzodiazepines for the relief of breathlessness in advanced malignant and non-malignant diseases in adults. *Cochrane Database Syst Rev*. 2016;10(10):Cd007354.
- Van Schayck OCP, Williams S, Barchilon V, Baxter N, Jawad N, Katsaounou PA, Kirenga BJ, Panaitescu C, Tsiligianni IG, Zwar N, Ostrem A. Treating tobacco dependence: guidance for primary care on life-saving interventions. Position statement of the IPCRG. *NPJ Prim Care Respir Med*. 2017 Jun 9;27(1):38. doi: 10.1038/s41533-017-0039-5. Erratum in: *NPJ Prim Care Respir Med*. 2017 Sep 5;27(1):52.
- Hardinge M, Annandale J, Bourne S, Cooper B, Evans A, Freeman D, et al. British Thoracic Society guidelines for home oxygen use in adults. *Thorax*. 2015;70 Suppl 1:i1-43.
- Abernethy AP, McDonald CF, Frith PA, Clark K, Herndon JE, 2nd, Marcello J, et al. Effect of palliative oxygen versus room air in relief of breathlessness in patients with refractory dyspnoea: a double-blind, randomised controlled trial. *Lancet (London, England)*. 2010;376(9743):784-93.
- Holland AE, Hill CJ, Jones AY, McDonald CF. Breathing exercises for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2012;10:Cd008250.
- Luckett T, Phillips J, Johnson MJ, Farquhar M, Swan F, Assen T, et al. Contributions of a hand-held fan to self-management of chronic breathlessness. *Eur Respir J*. 2017;50(2).